


# A new classification system for degenerative spondylolisthesis of the lumbar spine

Olivier Gille<sup>1</sup> · Houssam Bouloussa<sup>1</sup> · Simon Mazas<sup>1</sup> · Claudio Vergari<sup>2</sup> · Vincent Challier<sup>1</sup> · Jean-Marc Vital<sup>1</sup> · Pierre Coudert<sup>1</sup> · Soufiane Ghailane<sup>1</sup> 

Received: 16 January 2017 / Accepted: 17 August 2017  
© Springer-Verlag GmbH Germany 2017

## Abstract

**Purpose** There is no consensus for a comprehensive analysis of degenerative spondylolisthesis of the lumbar spine (DSLS). A new classification system for DSLS based on sagittal alignment was proposed. Its clinical relevance was explored.

**Methods** Health-related quality-of-life scales (HRQOLs) and clinical parameters were collected: SF-12, ODI, and low back and leg pain visual analog scales (BP-VAS, LP-VAS). Radiographic analysis included Meyerding grading and sagittal parameters: segmental lordosis (SL), L1–S1 lumbar lordosis (LL), T1–T12 thoracic kyphosis (TK), pelvic incidence (PI), pelvic tilt (PT), and sagittal vertical axis (SVA). Patients were classified according to three main types—1A: preserved LL and SL; 1B: preserved LL and reduced SL ( $\leq 5^\circ$ ); 2A: PI–LL  $\geq 10^\circ$  without pelvic compensation (PT  $< 25^\circ$ ); 2B: PI–LL  $\geq 10^\circ$  with pelvic compensation (PT  $\geq 25^\circ$ ); type 3: global sagittal malalignment (SVA  $\geq 40$  mm).

**Results** 166 patients (119 F: 47 M) suffering from DSLS were included. Mean age was  $67.1 \pm 11$  years. DSLS

demographics were, respectively: type 1A: 73 patients, type 1B: 3, type 2A: 8, type 2B: 22, and type 3: 60. Meyerding grading was: grade 1 ( $n = 124$ ); grade 2 ( $n = 24$ ). Affected levels were: L4–L5 ( $n = 121$ ), L3–L4 ( $n = 34$ ), L2–L3 ( $n = 6$ ), and L5–S1 ( $n = 5$ ). Mean sagittal parameter values were: PI:  $59.3^\circ \pm 11.9^\circ$ ; PT:  $24.3^\circ \pm 7.6^\circ$ ; SVA:  $29.1 \pm 42.2$  mm; SL:  $18.2^\circ \pm 8.1^\circ$ . DSLS types were correlated with age, ODI and SF-12 PCS ( $\rho = 0.34$ ,  $p < 0.05$ ;  $\rho = 0.33$ ,  $p < 0.05$ ;  $\rho = -0.20$ , and  $p = 0.01$ , respectively).

**Conclusion** This classification was consistent with age and HRQOLs and could be a preoperative assessment tool. Its therapeutic impact has yet to be validated.

**Level of evidence** 4.

**Keywords** Degenerative spondylolisthesis · Lumbar spine · Classification system · Spondylolisthesis · Clinical relevance

## Introduction

Degenerative spondylolisthesis of the lumbar spine (DSLS) is a common cause of consultation with spinal surgeons. Initially described by the obstetrician Herbiniaux [1], the term spondylolisthesis was first used by Kilian [2]. DSLS is thought to be caused by various degenerative processes affecting the intervertebral disc and facet joints responsible for the translation and slippage of one vertebral body onto the subjacent one. Its pathogenesis still remains unclear. DSLS typically occurs at the L4–L5 level in women older than 50 with a high pelvic incidence (PI) [3–5]. It is also frequently associated with spinal stenosis [6]. These degenerative modifications contribute to produce the following symptoms: lower back pain, leg pain, postural

---

Oral communication at the SICOT (International Society of Orthopaedic Surgery and Traumatology), annual meeting in Roma, Italy, September 8, 2016. Abstract no. 43858.  
Oral communication in IMAST annual meeting in Cape Town, South Africa, July 14, 2017.

---

✉ Soufiane Ghailane  
soufiane.ghailane@gmail.com

<sup>1</sup> Department of Spinal Surgery Unit 1, C.H.U Tripode Pellegrin, Bordeaux University Hospital, Université de Bordeaux, Place Amélie Raba Léon, 33076 Bordeaux, France

<sup>2</sup> School of Physics and Astronomy, University of Exeter, Exeter, UK

syndrome, and neurogenic claudication. Various classifications attempted to provide further understanding of this disease. However, they were based on etiology, topography, or slippage grading (percentage), and were restricted to a segmental analysis [7–10]. Therefore, the role of regional or global malalignment was not considered. None of these classification systems provide surgeons with a comprehensive analysis of DSLS or guidance for optimal care. Recently, several studies reported the close relationship between DSLS and sagittal alignment [11–14]. Spinopelvic malalignment plays a significant role in multiple spinal conditions [15–17]. It seems crucial to consider this parameter analyzing DSLS using preoperative full spine imaging.

A new classification system of DSLS based on sagittal alignment was proposed by Gille et al. [18]. The clinical relevance of this new classification system remains to be determined to confirm or not its clinical value. This aspect was addressed in the present study by analyzing the relationships between the different types of DSLS and patient demographics, radiographical parameters, and health-related quality-of-life scales (HRQOLs).

## Methods

### Study design and population

All patients admitted to our spinal surgery department for surgical treatment of DSLS with spinal stenosis (central, lateral recess, or foraminal) were retrospectively included between January 2011 and December 2015 following approval from our Institutional Review Board. The inclusion criteria were: (1) age >18 years; (2) degenerative spondylolisthesis of the lumbar spine requiring surgical treatment due to back pain associated with either neurogenic claudication or severe radiculopathy despite 6 months of optimal medical treatment and/or motor neurological deficit; (3) complete data (demographic information, health-related quality-of-life scales, and full standing spine X-rays).

Patients were excluded if they presented with: (1) a coronal malalignment with coronal Cobb angle >10°; (2) other causes of spondylolisthesis (isthmic, congenital, traumatic, and iatrogenic); (3) previous lumbar spine surgery; (4) active infection or neoplasm.

### Clinical parameters and health-related quality-of-life scales

Clinical parameters (age, gender, and body mass index) and health-related quality-of-life scales (HRQOLs) were collected: Short Form-12 Questionnaire (SF-12), Oswestry

Disability Index (ODI), and back pain and leg pain visual analog scales (BP-VAS and LP-VAS).

### Radiographical parameters

Radiographic analysis included slippage level, slippage percentage, Meyerding grading, and sagittal parameters: segmental lordosis (SL), L1–S1 lumbar lordosis (LL), T1–T12 thoracic kyphosis (TK), pelvic incidence (PI), pelvic tilt (PT), and sagittal vertical axis (SVA).

### The classification system

The proposed classification was based on the rating of sagittal full-body standing radiographs (EOS system, EOS imaging, Paris, France) used in routine. It was derived from the sagittal modifiers of the SRS-Schwab classification for adult spinal deformity (ASD) [19]. The SRS-Schwab classification for ASD was shown to be correlated with HRQOLs [20]. Two orthopedic surgeons performed all radiographical measurements for each patient using a validated software (Surgimap<sup>®</sup> Nemaris Inc., New York, NY, USA).

A formal description of the classification is given in Table 1; briefly, type 1 corresponds to a harmonious and aligned spine (Fig. 1), type 2 corresponds to a compensated spinal malalignment (Fig. 2), and type 3 corresponds to an altered global sagittal alignment (Fig. 3). Severity increases from type 1 to 3. Subtypes depend on segmental lordosis (type 1) or pelvic compensation (type 2). All patients were classified according to this classification system (Fig. 4).

### Statistical analysis

Differences of clinical or sagittal spinal parameters according to spondylolisthesis types were assessed with non-parametric Kruskal–Wallis tests. Correlations between demographic data, HRQOLs, and radiographical parameters were assessed using Spearman's rank test, while differences were assessed with Mann–Whitney tests. Statistical analyses were performed using Matlab 2015b (Mathworks, Natick, MA, USA); statistical significance was set at  $p = 0.05$ .

## Results

### Demographic data

A total of 166 patients who underwent surgery in our spinal surgery department with complete data were included. There were 119 females and 47 males with a mean age of

**Table 1** Description of the classification system and patients

Type	Description	Parameters	Sub-type	Sub-type description	Age	Number of patients (males/females)
Type 1	LL adapted to PI (harmonious spine)	PI-LL <10° (Fig. 1)	1A	Preserved segmental lordosis (SL)	64 ± 11	73 (22/51)
			1B	Altered SL, with preserved LL	62 ± 10	3 (2/1)
Type 2	Compensated malalignment	PI-LL >10° (Fig. 2)	2A	Preserved global alignment without pelvic compensation (Pelvic tilt PT < 25°)	65 ± 12	8 (4/4)
			2B	Preserved global alignment with pelvic compensation (PT > 25°)	66 ± 10	22 (11/21)
Type 3	Altered global alignment (SVA >40 mm)	SVA >40 mm (Fig. 3)	3		72 ± 9	60 (18/42)
Average					66 ± 10	166 (47/119)

67.1 ± 10.5 years at surgery. All patients had DSLS with spinal stenosis. The majority of patients in this study had neurogenic claudication due to central spinal stenosis (90%). The remaining 10% suffered from lateral recess or foraminal stenosis. Affected levels were: L4–L5 ( $n = 121$ ), L3–L4 ( $n = 34$ ), L2–L3 ( $n = 6$ ), and L5–S1 ( $n = 5$ ). DSLS classification demographics were, respectively: type 1A ( $n = 73$ ), type 1B ( $n = 3$ ), type 2a ( $n = 8$ ), type 2B ( $n = 22$ ), and type 3 ( $n = 60$ ). The mean BMI was  $26.14 \pm 5.05 \text{ kg/m}^2$ .

The demographic distribution of spondylolisthesis types is reported in Table 1.

### Preoperative clinical parameters

The mean LP-VAS and BP-VAS were, respectively,  $6 \pm 2$  and  $7 \pm 2$ ; pain did not correlate with any other parameter. The mean ODI was  $48 \pm 15$ . The mean SF-12 PCS was  $31 \pm 8$ .

Clinical parameter data classified by type are reported in Table 2.

### Radiographical parameters

The Meyerding grading was the following: grade 1 ( $n = 124$ ) and grade 2 ( $n = 24$ ). The mean slippage percentage was  $15 \pm 7.6\%$ . The mean values of spinopelvic parameters were: PI ( $59.3^\circ \pm 11.9^\circ$ ), PT ( $24.3^\circ \pm 7.6^\circ$ ), SS ( $35^\circ \pm 9^\circ$ ), PI-LL ( $9^\circ \pm 12^\circ$ ), SL ( $18.2^\circ \pm 8.1^\circ$ ), LL ( $51.3^\circ \pm 13.1^\circ$ ), TK ( $41.0^\circ \pm 13.9^\circ$ ), and SVA ( $29.1 \pm 42.2 \text{ mm}$ ).

Radiographical parameter values are reported in Table 3.


PI-LL was correlated with ODI ( $\rho = 0.24$ ,  $p = 0.002$ ). SVA was correlated with ODI ( $\rho = 0.3$ ,  $p = 0.0002$ ) and SF12-PCS ( $\rho = -0.18$ ,  $p = 0.02$ ). PT, PI, LL, TK, and SL were not correlated with HRQOLs.

### Relationships between spondylolisthesis types, and clinical and radiographical parameters

DSLS types were correlated with age, ODI, and SF-12 PCS ( $\rho = 0.34$ ,  $p < 0.05$ ;  $\rho = 0.33$ ,  $p < 0.05$ ;  $\rho = -0.20$ ,  $p = 0.01$ , respectively). Type 3 patients had a significantly higher ODI than type 1A and 1B patients ( $p = 0.0002$ ), while SF-12 PCS was significantly lower in type 3 than type 1A and 1B patients ( $p = 0.03$ ), demonstrating a quality-of-life degradation with increased type severity. BP-VAS and LP-VAS did not vary with types. Low SL did not influence HRQOLs.

Relationships between the spondylolisthesis classification, age, and HRQOLs are reported in Fig. 5 while effects of age on spinal parameters are detailed in Fig. 6. Trends were observed between segmental parameters (SL and slippage percentage) and classification types. SL decreased with increasing types, with a significant difference between type 1 and type 3 ( $p = 0.02$ , Fig. 6), while slippage percentage increased with increasing types, again with a significant difference between type 1 and type 3 ( $p = 0.01$ , Fig. 6). Furthermore, LL decreased with increasing types. Differences between type 1 and type 3 were statistically significant ( $p < 0.001$ , Fig. 6). Mean PI in type 1 patients was physiological ( $56.3^\circ \pm 9.4^\circ$ ), while it was significantly increased in type 2 ( $63.9^\circ \pm 12^\circ$ ,  $p < 0.001$ ) and type 3 ( $69^\circ \pm 13.7^\circ$ ,  $p < 0.001$ ). Mean PT was the highest in type 2 ( $29.7^\circ \pm 6.3^\circ$ ,  $p < 0.05$ ). SVA expectedly tended to increase with increasing types. Type 2 patients had significantly lower TK ( $32.3^\circ \pm 13.5^\circ$ ) than type 1 ( $43.9^\circ \pm 11.3^\circ$ ,  $p = 0.00002$ ) and type 3 patients ( $41.8^\circ \pm 15.4^\circ$ ,  $p = 0.004$ ).

Relationships between classification types and radiographical parameters are reported in Fig. 6.




**Fig. 1** Type 1 on lateral standing low-dose X-ray view. Harmonious and aligned spine

## Discussion

DSLS is a common cause of lower back pain and leg pain with or without neurogenic claudication. It is caused by several degenerative modifications [5]. The prevalence of patients with symptomatic DSLS is expected to rise as the population ages. The other known causes of spondylolisthesis (isthmic, congenital, traumatic, and iatrogenic) are not considered here.

To our knowledge, there is no classification system or tool providing surgeons with a comprehensive analysis of sagittal alignment in DSLS. In this framework, we proposed a new classification system based on sagittal alignment with three main types.

The management of DSLS requires a holistic and comprehensive analysis of each case. Recently, different studies reported that spinopelvic sagittal malalignment played an important role in multiple spinal conditions [15–17] and especially in the management of DSLS




**Fig. 2** Type 2 on lateral standing low-dose X-ray view. Altered global LL, compensated malalignment

[11, 12]. Standing lateral radiographs are the most appropriate, non-invasive test to detect degenerative DSLS [21]. However, the analysis of sagittal spinal alignment seems to be an important factor for the full assessment of DSLS [22]. Indeed, our own experience with the treatment of spondylolisthesis is that neglecting the role of sagittal alignment in DSLS, as shown by Kumar et al., may lead to poor clinical outcome and patient satisfaction [23]. We observed that patients mistreated as type 1 with a single-level posterior fusion, while they actually were type 2 or 3

required revision surgery to prolong constructs more frequently. However, this reflects our local experience and is not supported by clinical evidence; a longitudinal study is currently under way.

Our data were similar to the literature findings. Typically, the slippage was less than 30% [3–5, 24, 25]. In the present study, the sex ratio was 2.5. The mean age was  $67.1 \pm 11$  years. Mean slippage was  $14.6 \pm 7.6\%$ . Patients were older and the sex ratio was comparable to results from other studies. The mean PI was  $59.3^\circ \pm 11.9^\circ$ ,



**Fig. 3** Type 3 on lateral standing low-dose X-ray view. Altered global alignment

which is higher than in the general population ( $52.6^\circ \pm 10.4^\circ$  according to Mac Thiong et al. [26]) but comparable with other DSLS cohorts [18]. Indeed, patients with a high PI are predisposed to the development of DSLS [15, 27].

This classification system was consistent with age, ODI, and SF-12 PCS ( $\rho = 0.34$ ,  $p < 0.05$ ;  $\rho = 0.33$ ,  $p < 0.05$ ;  $\rho = -0.20$ , and  $p = 0.01$ , respectively). Aging is responsible for increasing clinical and radiographical DSLS severity and was, therefore, associated with increasing types. Indeed, compensatory mechanisms are progressively overrun due to muscular degeneration and osteoarthritis as


they become unable to restore sagittal imbalance. Furthermore, HRQOLs decreased with increasing types: the type definition was entirely based on X-ray measurements and had no direct link with age or HRQOL scores, which reduces the risk of bias. ODI and SF-12 PCS showed significant correlations, albeit weak, with several sagittal parameters (PI–LL correlated with ODI,  $p < 0.05$ , while SVA correlated with ODI and SF-12 PCS,  $p < 0.05$ ). It should be noted that low SL did not influence HRQOLs. This may be explained by the low number of patients with  $SL < 5^\circ$  (three patients). Classification types were not correlated with LP-VAS, BP-VAS, and SF-12 MCS. In fact, these parameters are known as highly subjective and their value was limited in the absence of a comparison with postoperative values.

Our results support the well-described natural history of DSLS featuring decreasing SL and disc height loss in parallel with increasing slippage. Indeed, type 3 presented a lower SL and LL compared to type 1 ( $p = 0.02$ ) with a linear decreasing trend. The slipping percentage significantly increased with types. Furthermore, PT was increased in types 2 and 3 compared to type 1. This increase was predominant in type 2, in accordance with the classification definition. This may be explained by overrun compensatory mechanisms in type 3. Type 2 patients managed to keep a “subnormal” SVA ( $< 40$  mm) by a PT increase. PT is the key of pelvic adaptation [28].

We hypothesize that there is a dynamic continuum from type 1 to type 3, which is yet to be proven following the same patients over time. Type 1A corresponds to balanced spines with preserved local and global sagittal balance. Type 1B includes a local compensation with disc flexion and loss of segmental lordosis. Type 2A and 2B include a PI–LL mismatch. This is due to multi-segmental degenerative disc disease responsible for a loss of LL. Type 2A (PI =  $57^\circ$ ) presented a lower PI than type 2B (PI =  $67^\circ$ ); in that sense, in these unbalanced but compensated subgroups, type 2A grossly corresponded to a flat lumbar spine with a mainly thoracic adaptation in hypokyphosis and type 2B corresponded to a dynamic lumbar spine with overrun thoracic adaptation in hypokyphosis and a mainly pelvic adaptation in retroversion. Indeed, patients in type 2 group displayed significantly lower thoracic kyphosis ( $32.3^\circ \pm 13.5^\circ$ ) than patients in type 1 ( $43.9^\circ \pm 11.3^\circ$ ) and type 3 ( $41.8^\circ \pm 15.4^\circ$ ). These compensatory mechanisms display specific limits with aging (thoracic extension muscular fatigability; pelvic maximum retroversion impaired by hip osteoarthritis). Type 3 represents a significant global malalignment resulting from overrun local and regional compensatory mechanisms (thoracic and pelvic); patients, therefore, commonly use walking canes.

Several authors extensively described the biomechanics of DSLS [11, 12, 15]. It appeared that patients with a

**Fig. 4** Decision-tree algorithm



**Table 2** Clinical characteristics by type

	LP-VAS	BP-VAS	ODI	SF-12 PCS
Type 1A	6 ± 2	7 ± 2	0.42 ± 0.12	39 ± 11
Type 1B	5 ± 1	6 ± 1	0.39 ± 0.14	42 ± 1
Type 2A	7 ± 1	8 ± 1	0.49 ± 0.17	35 ± 12
Type 2B	6 ± 2	7 ± 2	0.45 ± 0.16	37 ± 10
Type 3	6 ± 2	6 ± 2	0.52 ± 0.15	39 ± 10
Mean ± SD	6 ± 2	7 ± 2	0.46 ± 0.15	31 ± 8

Quality-of-life assessments: Short Form-12 Questionnaire Physical Composite Scale (SF-12 PCS), Oswestry Disability Index (ODI), and Low Back and Leg Pain Visual Analog Scale (BP-VAS; LP-VAS)

dynamic lumbar spine and high PI (Roussouly 3 or 4) were prone to developing slippage [13, 29]. A high PI is, therefore, the initial driving force behind the development of DSLS, supported by the high mean PI observed in types 2 and 3 (Fig. 6). Thus, we believe that degenerative disc disease occurs later in the natural history of DSLS, resulting either in single-level disc degeneration with local kyphosis (type 1B) or multi-level disc degeneration with global hypolordosis (type 2). Type 1B seemed to be an


isolated and rare (three patients) entity due to its lower PI compared with all other types (PI = 45° ± 9°). This may be explained by the very nature of the process behind the development of DSLS. Indeed, static spines (Roussouly 1 and 2) with low PI and low SS are less frequently responsible for DSLS. Type 2A only represented 4.8% of the population in the present study. Indeed, according to Liu et al. [15], increased lumbar lordosis and pelvic incidence account for the high sheering forces responsible for the development of spondylolisthesis. This may explain the higher prevalence of type 2B (13.25%) over type 2A (4.8%). Type 3 represents the final stage of DSLS and occurs in significantly older patients (72 ± 9 years, *p* = 0.006).

One of the limitations of this study was the absence of evaluation of the impact of spinal stenosis on posture. MRI analysis seems mandatory and could prevent to over treat type 2, as deformity cases. Discerning the implication of spinal stenosis on posture would require a control group with no spinal stenosis. However, all patients suffered from neurological symptoms and if postural factors intervened, the distribution of postural factors was also assumed uniform


**Table 3** Radiographical characteristics by types

	PI (°)	LL (°)	PI-LL (°)	PT (°)	SS (°)	SL (°)	SVA (mm)	TK (°)	Slippage (%)
Type 1A	57 ± 9	57 ± 9	1 ± 6	21 ± 5	36 ± 8	21 ± 7	11 ± 17	44 ± 11	13 ± 8
Type 1B	45 ± 9	40 ± 9	6 ± 1	17 ± 5	28 ± 4	3 ± 1	13 ± 4	36 ± 15	12 ± 7
Type 2A	57 ± 11	46 ± 17	11 ± 10	22 ± 1	35 ± 11	19 ± 11	18 ± 15	30 ± 21	12 ± 6
Type 2B	67 ± 12	49 ± 14	18 ± 6	33 ± 5	34 ± 10	16 ± 8	14 ± 17	33 ± 10	16 ± 6
Type 3	61 ± 14	47 ± 15	16 ± 13	26 ± 9	35 ± 11	16 ± 8	73 ± 34	42 ± 15	17 ± 8
Mean ± SD	59 ± 12	51 ± 13	9 ± 12	24 ± 8	35 ± 9	18 ± 8	34 ± 38	41 ± 14	15 ± 8

Spinal parameters: pelvic incidence (PI), lumbar lordosis (LL), pelvic tilt (PT), sacral slope (SS), segmental lordosis (SL), sagittal vertical axis (SVA), and thoracic kyphosis (TK)



**Fig. 5** Relationship between degenerative spondylolisthesis types, age, and quality-of-life indexes. *ODI* Oswestry Disability Index, *SF-12 PCS* Short Form-12 Questionnaire Physical Composite Scale



**Fig. 6** Relationships between degenerative spondylolisthesis types and radiographical spinal characteristics: Pelvic incidence (*PI*), pelvic tilt (*PT*), segmental lordosis (*SL*), sagittal vertical axis (*SVA*), L1–S1 lumbar lordosis (*LL*), and slippage

in the studied population, since there was no control group without spinal stenosis. Since the surgical treatment of DSLs with isolated back pain and no symptomatic spinal stenosis (no neurogenic claudication or radiculopathy) remains highly controversial, we believe that this classification should not be used in such cases. Spinal stenosis has been demonstrated to be a cause of reversible lumbar kyphosis [30]. Buckland et al. studied different posture patterns between patients with either ASD or degenerative lumbar stenosis (DLS), and concluded that they engaged different compensatory mechanisms [30]. Indeed, according to

Buckland et al., patients attempt to decompress neural elements by permitting truncal sagittal malalignment driven by a posterior pelvic shift. The latter was recruited earlier in patients with DLS compared with patients suffering from ASD. Besides, patients in mild-to-moderate malalignment did not recruit PT until moderate-to-severe malalignment was present. They also showed that increasing SVA before recruiting PT was the preferable mode of compensation for patients with DLS. Those fundamental differences in terms of compensation behavior advocate the use of this classification for DSLs with spinal stenosis only.



Considering solely PI and other sagittal lumbar parameters for an optimal surgical management of DSLS occults fundamental regional and global dynamic compensatory mechanisms. This classification proposes to fully integrate sagittal spinal balance and pelvic parameters taking into account commonly used preoperative criteria. Furthermore, Smith et al. showed that patients with improved spinopelvic sagittal modifiers (PI–LL, PT, or SVA) after surgical correction had significantly higher HRQOLs than those whose modifiers deteriorated or remained the same [29]. However, the therapeutic impact of this classification has to be validated.

A therapeutic guidance can be proposed according to this classification. The ideal goal of surgical management is to maintain, approach, or restore a physiological postoperative spinal balance:

- Type 1: a segmental approach is advised: decompression and fusion alone with no correction or dynamic stabilization (type 1A) [31]. Type 1B includes segmental kyphosis ( $SL < 5^\circ$ ) and we believe that it is preferable in that case to restore SL using an intersomatic device, through an anterior or posterior approach.

Treatments for type 2 and 3 are similar to strategies developed for ASD: regional correction becomes essential to reach a satisfying postoperative global alignment.

- Type 2: there is PI–LL mismatch. Patients compensate with thoracic spine extension (flat back appearance) (Type 2A,  $PT < 25^\circ$ ) or with pelvic retroversion (type 2B,  $PT > 25^\circ$ ). The aim in these cases is to restore a harmonious spine with an LL adapted to PI.
- Type 3: sagittal imbalance prevails ( $SVA > 40$  mm). More aggressive surgical treatment may be considered to correct sagittal malalignment, especially in case of significant clinical sagittal imbalance. Treating only the slippage level may lead to a poor clinical outcome.

## Conclusion

This classification fully combines segmental, regional, and global analysis of sagittal balance with regard to DSLS. Classification types were consistent with age and HRQOLs (ODI, SF12-PCS). This classification potentially represents a useful tool for comprehensive analysis of DSLS before surgical treatment taking into account sagittal balance. Further clinical evidence is currently being collected to validate its therapeutic impact.

## Compliance with ethical standards

**Conflict of interest** No funds were received in support of this work. No benefits in any forms have been or will be received from a

commercial party related directly or indirectly to the subject of this manuscript.

## References

1. Herbiniaux G (1782) *Traité sur Diverse Accouchements Laborieux et sur les Polypes de la Matrice*
2. Kilian HF (1854) *Schilderungen neuer Beckenformen und ihres Verhaltens im Leben*. Bassermann & Mathy
3. Jacobsen S, Sonne-Holm S, Røvsing H et al (2007) Degenerative lumbar spondylolisthesis: an epidemiological perspective: the Copenhagen osteoarthritis study. *Spine* 32:120–125
4. Love TW, Fagan AB, Fraser RD (1999) Degenerative spondylolisthesis. *Bone Jt J* 81:670–674
5. Matsunaga S, Sakou T, Morizono Y et al (1990) Natural history of degenerative spondylolisthesis: pathogenesis and natural course of the slippage. *Spine* 15:1204–1210
6. Iguchi T, Wakami T, Kurihara A et al (2002) Lumbar multilevel degenerative spondylolisthesis: radiological evaluation and factors related to anterolisthesis and retrolisthesis. *Clin Spine Surg* 15:93–99
7. Kepler CK, Hilibrand AS, Sayadipour A et al (2015) Clinical and radiographic degenerative spondylolisthesis (CARDS) classification. *Spine J* 15:1804–1811
8. MacNab I (1950) Spondylolisthesis with an intact neural arch—the so-called pseudo-spondylolisthesis. *Bone Jt J* 32:325–333
9. Meyerding HW (1932) Spondylolisthesis: surgical treatment and results. *Surg Gynecol Obstet* 54:371–377
10. Wiltse LL, Newman PH, Macnab IAN (1976) Classification of spondyloistis and spondylolisthesis. *Clin Orthop* 117:23–29
11. Funao H, Tsuji T, Hosogane N et al (2012) Comparative study of spinopelvic sagittal alignment between patients with and without degenerative spondylolisthesis. *Eur Spine J* 21:2181–2187
12. Kim MK, Lee S-H, Kim E-S et al (2011) The impact of sagittal balance on clinical results after posterior interbody fusion for patients with degenerative spondylolisthesis: a pilot study. *BMC Musculoskelet Disord* 12:1
13. Roussouly P, Pinheiro-Franco JL (2011) Biomechanical analysis of the spino-pelvic organization and adaptation in pathology. *Eur Spine J* 20:609–618
14. Schwab FJ, Blondel B, Bess S et al (2013) Radiographical spinopelvic parameters and disability in the setting of adult spinal deformity: a prospective multicenter analysis. *Spine* 38:E803–E812
15. Liu H, Li S, Zheng Z et al (2015) Pelvic retroversion is the key protective mechanism of L4–5 degenerative spondylolisthesis. *Eur Spine J* 24:1204–1211
16. Protosaltis T, Schwab F, Bronsard N et al (2014) The T1 pelvic angle, a novel radiographic measure of global sagittal deformity, accounts for both spinal inclination and pelvic tilt and correlates with health-related quality of life. *J Bone Jt Surg Am* 96:1631–1640
17. Schwab F, Patel A, Ungar B et al (2010) Adult spinal deformity—postoperative standing imbalance: how much can you tolerate? An overview of key parameters in assessing alignment and planning corrective surgery. *Spine* 35:2224–2231
18. Gille O, Challier V, Parent H et al (2014) Degenerative lumbar spondylolisthesis. Cohort of 670 patients, and proposal of a new classification. *Orthop Traumatol Surg Res* 100:S311–S315
19. Lowe T, Berven SH, Schwab FJ, Bridwell KH (2006) The SRS classification for adult spinal deformity: building on the King/Moe and Lenke classification systems. *Spine* 31:S119–S125
20. Terran J, Schwab F, Shaffrey CI et al (2013) The SRS-Schwab adult spinal deformity classification: assessment and clinical

- correlations based on a prospective operative and nonoperative cohort. *Neurosurgery* 73:559–568
21. Matz PG, Meagher RJ, Lamer T et al (2016) Guideline summary review: an evidence-based clinical guideline for the diagnosis and treatment of degenerative lumbar spondylolisthesis. *Spine J* 16:439–448
  22. Challier V, Boissiere L, Obeid I et al (2017) One-level lumbar degenerative spondylolisthesis and posterior approach: is transforaminal lateral interbody fusion mandatory?: a randomized controlled trial with 2-year follow-up. *Spine* 42:531–539
  23. Kumar M, Baklanov A, Chopin D (2001) Correlation between sagittal plane changes and adjacent segment degeneration following lumbar spine fusion. *Eur Spine J* 10:314–319
  24. Cummins J, Lurie JD, Tosteson T et al (2006) Descriptive epidemiology and prior healthcare utilization of patients in the spine patient outcomes research trial's (sport) three observational cohorts: disc herniation, spinal stenosis and degenerative spondylolisthesis. *Spine* 31:806
  25. Rosenberg NJ (1975) Degenerative spondylolisthesis. Predisposing factors. *J Bone Jt Surg Am* 57:467–474
  26. Mac-Thiong J-M, Roussouly P, Berthonnaud É, Guigui P (2010) Sagittal parameters of global spinal balance: normative values from a prospective cohort of seven hundred nine Caucasian asymptomatic adults. *Spine* 35:E1193–E1198
  27. Barrey C, Jund J, Nosedà O, Roussouly P (2007) Sagittal balance of the pelvis-spine complex and lumbar degenerative diseases. A comparative study about 85 cases. *Eur Spine J* 16:1459–1467
  28. Lafage V, Schwab F, Patel A et al (2009) Pelvic tilt and truncal inclination: two key radiographic parameters in the setting of adults with spinal deformity. *Spine* 34:E599–E606
  29. Roussouly P, Gollogly S, Berthonnaud E, Dimnet J (2005) Classification of the normal variation in the sagittal alignment of the human lumbar spine and pelvis in the standing position. *Spine* 30:346–353
  30. Buckland AJ, Vira S, Oren JH et al (2016) When is compensation for lumbar spinal stenosis a clinical sagittal plane deformity? *Spine J* 16:971–981
  31. Liang H-F, Liu S-H, Chen Z-X, Fei Q-M (2017) Decompression plus fusion versus decompression alone for degenerative lumbar spondylolisthesis: a systematic review and meta-analysis. *Eur Spine J*. doi:[10.1007/s00586-017-5200-x](https://doi.org/10.1007/s00586-017-5200-x)